

# Health Information System

Organisation: \_\_\_\_\_

## 10.0 HIV/AIDS

Location: \_\_\_\_\_

### Emergency Reporting Pad

For use in emergency settings





# Health Information System

Emergency Reporting Form

Organisation: \_\_\_\_\_

Location: \_\_\_\_\_

Reporting period: \_\_\_\_\_

## 10.0 HIV/AIDS

### 10.1 Condom Distribution

Number of male condoms distributed	
------------------------------------	--

### 10.2 HIV Testing and Counselling (HCT)

Number of HCT clients	< 18		≥ 18	
	M	F	M	F
Pre-test counselled				
Tested for HIV				
Tested positive for HIV				
Post-test counselled				

### 10.3 PMTCT (Antenatal)

Number of pregnant women	< 18	≥ 18	Partner
Pre-test counselled			
Tested for HIV			
Tested positive for HIV			
Post-test counselled			
Who accepted to take ARV at 28 weeks			

## 10.4 PMTCT (Labour and Delivery)

HIV positive deliveries	< 18		≥ 18	
	Home	Health Centre	Home	Health Centre
Live births				
Still births				
Abortions				
During which mother swallowed ARV				
After which newborn was given ARV < 72 hours				
No. of mother-newborn pairs that received ARV *				

\* on time, according to national protocol

## 10.5 PMTCT (Postnatal)

Number of HIV positive women who:		< 18	≥ 18	Total
Choose to exclusively breastfeed				
Choose to replacement feed				
Received at least 1 HBC visit*				
Accepted modern family planning				
Number of mothers who started co-trimoxazole				
Number of infants who started co-trimoxazole				
Number of exits:	<i>discharge</i>			
	<i>death (neonatal)</i>			
	<i>death (&lt; 1 year)</i>			
	<i>death (&gt; 1 year)</i>			
	<i>default</i>			
	<i>referral</i>			
Infant HIV outcomes	HIV positive at 18 months			

\* Home Based Care