


Helping Mothers and Babies Survive Pre-Eclampsia & Eclampsia

ACTION PLAN 2 - ADVANCED CARE


SEVERE PRE-ECLAMPSIA

Pre-Eclampsia and
One or More Danger Signs



ECLAMPSIA

Convulsions or Unconscious



Receive referral OR Continue care

Begin OR Continue MgSO₄


Begin OR Continue medication
to reduce severe BP

Continue close monitoring
of woman and fetus

Within
4 hours

If no convulsions and
conscious

If convulsions or
unconscious



Confirm Gestational Age

Deliver

Continue MgSO₄
for 24 hours after birth
or last convulsion

Within
12 hours

Pre-viable

Viable to < 34 weeks

≥ 34 week to
< 37 weeks

> 37 weeks

Within
24 hours

End pregnancy



Give
dexamethasone

Within
24 hours

Continue MgSO₄
for 24 hours after birth
or last convulsion

Admit woman and
monitor closely:

- Convulsions
- Uncontrolled severe hypertension
- Worsening maternal condition (danger signs, exam, lab tests)
- Worsening fetal condition or demise
- Reaches 37 weeks 0 days

Yes to any

Continue medication
to reduce severe BP



Provide essential
maternal and
newborn care

Continue to monitor
after birth